## **SOFTBALL** SAFETY

A PARENT'S GUIDE FOR KEEPING KIDS IN THE GAME

ENS INTE

Softball is a non-contact sport so the risk of injury is much lower than other sports. However, shoulder, knee, and ankle injuries are common. This reference guide provides information on softball injuries that require treatment.

## SHOULDER INJURIES

Shoulder injuries are common injuries in softball. They most often are caused by the collision that is forced when one player drives another into the boards or ice. Seek immediate medical care when the collar bone appears deformed or if the athlete indicates the shoulder is "out of socket." Symptoms include:

- Pain when performing the activity or sport
- Dull pain even at rest
- Intermittent swelling
- Loss of motion at the shoulder or wrist
- Decreasing performance

### **ANKLE INJURIES**

The most common injury in sports is a **lateral ankle sprain**, which occurs by rolling the ankle over the outside of the foot. A lateral ankle sprain causes damage to the ligaments just below the bone on the outside of the ankle. In some cases a "pop" is felt or heard by the athlete. Treatment varies with the severity of the injury:

- Mild sprains require rest but not necessarily medical treatment.
- Injuries with persistent swelling, pain or any deformity should be seen by a physician.

### **HEAD INJURIES**

A **concussion** is a brain injury usually caused by a sudden jolt or a blow to the head or neck that disrupts normal brain function. An athlete does not need to be knocked out or have memory loss to have suffered a concussion. You may observe that an athlete with a concussion:

- Appears dazed or stunned
- Moves clumsily
- Answers questions slowly
- Forgets plays
- Has behavior or personality changes
- Is unsure of game, score or opponent
- Can't recall events either before or after hit
- Loses consciousness
- An athlete with a concussion may have:
- Headache
- Concentration or memory problems
- Nausea
- Double or fuzzy vision
- Balance problems or dizziness
- Feelings of being "in a fog"
- Sensitivity to light or noise

An athlete with signs of a concussion should be removed from play immediately and not allowed to return until evaluated by a doctor. Do not leave an athlete alone after a concussion.

#### DEHYDRATION

Softball players are at risk of **dehydration** if they don't get enough fluid to replace what is lost through the skin as sweat and through the lungs while breathing. It is important to drink plenty of fluids before, during and after a workout or game. An athlete's performance can be impacted by even mild dehydration. Athletes should take a water bottle to school and drink between classes and during breaks so that they are well-hydrated before their workout. In addition:

- Water should be readily available when working out.
- Athletes should drink often, ideally every 15 to 30 minutes.
- Sports drinks are recommended for activities lasting longer than one hour to replace sugar and salt as well as water.

Early signs of dehydration can be non-specific and include fatigue, nausea, decreased athletic performance, headache, apathy, irritability and thirst. Signs of advanced dehydration include dark urine, decrease in reaction time, dry lips and mouth and disorientation.



# SOFTBALL SAFETY (CONT.)

### SPORTS SAFETY

Children ages 5 to 14 make up almost 40 percent of all sports injuries treated in hospital emergency rooms. Injuries in children are best handled by pediatric specialists trained in treating skeletally immature patients.

#### Athletes should see a pediatrician or pediatric sports medicine physician if any of these symptoms are present:

- Deformity
- Limping that lasts more than 48 hours
- Soft tissue swelling that gets worse the next day despite ice and over-the counter anti-inflammatory medication.
- Effusion mobile soft tissue swelling on both sides of a joint, often easily seen at the knee or ankle
- Pain that returns quickly with activity at the next session or is not gone after two weeks of rest.

#### How to protect your child:

- Schedule your child for an annual physical before playing sports.
- Monitor play and practice and encourage players to abide by the rules.
- Have a first aid kit handy and an emergency action plan in place.
- Keep sports fun! Remember to be positive and don't push kids to perform beyond their abilities.

## SSM HEALTH CARDINAL GLENNON SPORTSCARE

SSM Health Cardinal Glennon SportsCare is the premier pediatric sports medicine provider in St. Louis and St. Charles. We provide exclusive, direct access to comprehensive medical care for young athletes. By working with multiple specialists and care partners, we guarantee your child will get the top care that is best suited to treat their unique injury, improving recovery time and outcome.

Specialists your child has access to through SSM Health Cardinal Glennon SportsCare include orthopedists, pulmonologists, cardiologists, radiologists, neurologists, emergency medicine physicians, adolescent medicine physicians, rehabilitation specialists, physical therapists, nutritionists and pediatric psychologists.

Partners your child has access to through SSM Health Cardinal Glennon SportsCare include SSM Health Cardinal Glennon Children's Hospital, SSM Health Orthopedics, SSM Physical Therapy and SLU*Care* Physician Group of Saint Louis University. We also keep kids in the game through educational programs and support for parents, coaches and athletes that focus on injury prevention, proper technique and overall athletic health.

For more information about SSM Health Cardinal Glennon SportsCare or to find a specialist for your athlete, call us at **314-577-5640** or visit us at **cardinalglennon.com/sportscare**.

Expert care for young athletes by SSM Health Cardinal Glennon Children's Hospital and SSM Health emergency medicine specialists, **24 hours a day in the ER** at: SSM Health DePaul Hospital, SSM Health St. Clare Hospital, SSM Health St. Joseph Hospital - Lake Saint Louis, SSM Health St. Joseph Hospital - St. Charles.

Web

You

f



cardinalglennon.com/**sportscare** youtube.com/ssmcardinalglennon facebook.com/cardinalglennon twitter.com/cardinalglennon

Bumps, Bruises, Twists & Muscle Strains

These can affect all areas of the body.

Recommended treatment is the **PRICE** formula:

Protect the area with a sling or crutches, if necessary. Rest the injured area.

Ice the injury for 20 minutes at a time. Do not apply the ice directly to the skin.

 $C_{\mbox{ompress}}$  the injured area with a wrap. Do not pull tightly, as this can cut off circulation.

Elevate the injured area above the heart, if possible.

#### Make sure your young athlete:

- Wears appropriate properly-fitting safety gear, free of heavy wear and tear
- Stays properly hydrated. Kids don't sweat as much as adults and need to drink plenty of fluids before, during and after activity.
- Does warm-up and cool-down exercises before and after practices and games
- Gets proper rest and avoids overdoing it.